

## CHEMICAL COMPOSITION

Humic Acids	12 - 15 g/l
Fulvic Acids	8 - 10 g/l
Water-Soluble Humates	6 g/l
Micro Elements	Co, Mn, Mg, Cu, Zn, Fe
Macro Elements	P <sub>2</sub> O <sub>5</sub> , K <sub>2</sub> O, CaO, MgO, Na <sub>2</sub> O
18 Amino Acids	Asparagine, Threonine, Serine, Glutamine, Valine, Glycine, Leucine, Isoleucine, Phenylalanine, Methionine, Proline, Tyrosine, Lysine, Alanine, Histidine, Arginine
Vitamins	A, B1, B2, B3, B5, B6, B9, B12, E, C, D, P

## PHYSICAL PROPERTIES

Mass Fraction of Organic Matter	≥ 85%
Density	≥ 1.05 - 1.1 (g/cm <sup>3</sup> )
Moisture Content	85 - 90%
pH	7.5 - 9.5
State	Liquid

Available in: 1L Bottle, 5L Can, 1000L IBC  
Expiration Date: 12 months after opening



SCAN to LEARN  
MORE



SCAN to CHECK OUT  
DOSAGES



Healthy  
People



Natural  
Farming



Livestock



Poultry



Breeding



House Pets

**PRECOGROW** is a natural humic peat-sapropel concentrate, designed for livestock, farm poultry, breeding animals and pets.  
**Accelerate growth with our multipurpose feed additive PRECOCARE!**



UNIQUE PEAT-SAPROPEL FORMULA

NATURAL LIQUID  
HUMIC FEED ADDITIVE

PRECOCARE

HEADOFFICE: PRECOFERT s.r.o., ID No: 08097232  
Štítného 585/28, Prague 3, 130 00, Czech Republic  
+420 775 760 015 | [prodej@precofert.cz](mailto:prodej@precofert.cz)  
[eshop.precofert.cz](http://eshop.precofert.cz)

Designed for Immunity Boost,  
Metabolic Regulation and  
Growth Acceleration



# PRECOCARE

Our **PRECOCARE Concentrate** is composed of 70% Peat and 30% Sapropel. This proportion ensures the optimal balance of **humic** and **fulvic acids**, as well as other **essential nutrients** for livestock and farm poultry.

The ingredients are sourced from **clean and natural locations**, such as mountain lakes and pristine deposits, far from any industrial or urban pollution. The products consist solely of organic and eco-friendly elements: lake sapropel, low-moor peat, and lake water.

**A NATURAL TREASURE!**



## UNIQUE

Significantly increases the rate of **weight gain**, **immunity**, **metabolism**, **vitality** and **health**.



## SAFE

Safe for **both animals** and **humans**, with no allergenic or carcinogenic properties.



## COST-EFFICIENT

Increases the overall **feeding-efficiency** of livestock and poultry by **up to 20%** under the same feed, labor and environmental conditions.

## 70% + 30% PEAT - SAPROPEL FORMULA

### Humic & Fulvic Acids + Soluble Humates + Vitamins + Minerals + Amino Acids

Achieve a boost of **5-12%** in live weight, reduce mortality by **13-18%**, and lower feed consumption by **5-10%** per unit of weight gain.

### Compatability with Other Feed Additives

**PRECOCARE** can be added to feed or drinking water. It is compatible with other feed additives and has no effect on feed taste or digestibility.

- ✓ Dietary Enrichment
- ✓ Element Deficiency Prevention
- ✓ Regulation of Metabolic Processes
- ✓ Body Detoxification & Immunity Boost
- ✓ Animal Growth Acceleration
- ✓ Development of Beneficial Gastrointestinal Microflora

## USAGE & APPLICATION DOSAGES

1

### Mixing with Dry Feed Ingredients

PRECOCARE can be mixed directly with all types of dry feed ingredients on farms during daily feed preparation in feed mixers or feeding systems for (dairy) cattle.

2

### Spraying on the Surface of Ready-Made Industrial Pelleted, Extruded, or Granulated Feed

This method enhances the palatability and attractiveness of the feed while preventing dust formation. It can be employed by both feed manufacturers and farmers, depending on the availability and convenience of the equipment.

3

### Dissolving and Adding to Drinking Water

This method reduces odor and ammonia concentration in animal housing, improving air quality. It is applicable to farm animals kept on bedding or litter, including poultry, pigs, or rabbits.

4

### Soaking Feed in PRECOCARE Before Feeding

This method increases the solubility and bioavailability of PRECOCARE in the feed, enhancing its absorption by the animals. It is suitable for farmers who prepare their own feed on the farm, as it does not require any special equipment or processing.

TYPE AND AGE OF ANIMAL	DAILY DOSAGE PER 1 KG OF LIVE WEIGHT	APPLICATION / INTERRUPTION
Cows 30 days before calving	0.5 ml with food / drink	15 days / 10 days
Milking cows	0.5 ml with food, silage	15 days / 10 days
Calves 2-6 months	0.5 ml with food / drink	15 days / 10 days
Calves from 10 days 2 months	0.4 ml with skimmed milk	60 days
Piglets suckling mother's milk	0.4 ml with milk, porridge	15 days / 10 days
Weaned piglets	0.5 ml with food / drink	15 days / 10 days
Pregnant sow	0.5 ml with food / drink	Up to 60 days
Pregnant sheep and goats	0.5 ml with food / drink	15 days / 10 days
Lambs, kids up to 4 months	0.4 ml with food / drink	15 days / 10 days
Goats 4-8 months	0.5 ml with food / drink	15 days / 10 days
Horses, camels	2 ml with food / drink	28 days / 28 days
Rabbits	0.4 ml with food / drink	15 days / 10 days
Broiler chickens	2 l per 1000 l of water	The whole period
Laying hens	2 l per 1000 l of water	30 days / 7 days
Turkey, ducks, geese, ostriches	2 l per 1000 l of water	30 days / 7 days

**Have questions?** Please contact our support to find out more about **PRECOCARE**:

[info@precofert.cz](mailto:info@precofert.cz)

**PRECOCARE**  
Investing in Nature